arXiv:1211.2728v1 [hep-th] 12 Nov 2012

Preprint typeset in JHEP style - HYPER VERSION

Rational conformal field theory and matrix level for strings on a torus

Ali Nassar and Mark A. Walton

Physics and Astronomy Department, University of Lethbridge, Lethbridge, Alberta, Canada T1K 3M4
`nassar@uleth.ca`, `walton@uleth.ca`

Abstract: Gukov and Vafa gave a simple, geometric criterion for rationality of a conformal σ-model with a two-dimensional torus as a target space. The modular parameter τ and Kähler parameter ρ must take special values in an imaginary quadratic number field so that the torus and its mirror possesses the property of complex multiplication. On the other hand, Gannon has classified the modular invariant partition functions of the algebras $U_{m,K}$: Abeliam current algebras of the kind realized on a torus, but with a matrix-valued level K. We investigate the relation between the Gukov-Vafa geometric characterization of rationality and the algebraic results of Gannon. The geometric interpretation of the $U_{m,K}$ will be given for $m = 2$. The Gauss product is used to give a geometric interpretation of $U_{2,K}$ in terms of rational points in the Narain moduli space which correspond to complex multiplication tori. We also show that rational points on the Grassmannian of self-dual lattices correspond to complex multiplication tori. This is another way to show that the set of rational conformal field theories is dense in the Narain moduli space.

Contents

1. Introduction 1

2. Matrix-level Kac-Moody algebras 2

3. Strings on a complex multiplication torus 6

4. Rational points on Grassmannians and CM tori 11

5. Conclusion 13

1. Introduction

Two-dimensional conformal field theories (CFTs) are quite well understood. The local conformal symmetry described by the Virasoro algebra and its extensions is in many cases powerful enough to give a complete determination of the operator spectrum, as well as explicit formulae for the correlation functions. These constitute a complete non-perturbative solution of the theory (see [1, 2] for an introduction). Such theories give concrete instances of non-trivial fixed points of the renormalisation group, many of which have a realisation in statistical mechanical systems. They are also the building blocks of perturbative string theory where they describe the dynamical variables of the theory on the string world sheet.

A particularly simple class are *rational* conformal field theories (RCFTs), characterized by having a finite number of primary fields [3]. They are consistently described by a finite set $\mathcal{I}$ of representations $\mathcal{V}_i$, $i \in \mathcal{I}$, of a certain chiral algebra $\mathcal{A}$. Moreover the corresponding genus one-characters $\chi_i(q)$ form a finite dimensional unitary representation of the modular group $SL(2,\mathbb{Z})$

$$\chi_i\left(\frac{a\tau+b}{c\tau+d}\right) = \sum_j M_i^j \chi_j(\tau). \tag{1.1}$$

One arena where RCFTs arise is in string compactifications on Calabi-Yau manifolds at special values of their moduli, e.g., Gepner models [4]. It is important to understand the conditions for rationality in models which have continuously varying moduli. For example, the simplest compactification of a string theory are on tori– when are they described by rational conformal field theories? This question has been answered for the special case of a torus target in [5, 6, 7, 8] where a simple, geometric criterion for rationality was given. For T^2, the modular parameter τ and Kähler parameter ρ must take special values in the imaginary quadratic number field. A detailed study of RCFTs corresponding to T^2 target was performed in [8].

To see how a CFT can become rational at certain points of its moduli space, we revisit the simplest example of an RCFT: the $c = 1$ CFT of a compact free boson on a circle of rational square raduis $R^2 = p/q$. The radius R plays the role of a geometric modulus. At a generic value of R, the number of primary fields is infinite. At $R^2 = p/q$, the chiral algebra of this theory is enlarged. The infinite list of representations reconstitute themselves into finitely many irreducible representations of the enlarged chiral algebra. In the present case the chiral algebra is the Kac-Moody algebra generated by a $U(1)_k$ current $J(z) = i\sqrt{2k}\partial X$ extended by including charged fields $E^{\pm}(z) = \exp[\pm i\sqrt{2k}X(z)]$ of dimension k and charge $\pm 2k$, where $k = pq$. The representations of $U(1)_k$ which are local with respect to all vertex operators are labelled by an integer defined modulo $2k$. Defining the even integral lattice $\Gamma_k = \mathbb{Z}\sqrt{2k}$, the extended chiral algebra is the chiral vertex operator algebra $\mathcal{A}\big(\Gamma_k\big) = \langle J(z), E^{\pm}(z)\rangle$. The representations of the chiral algebra $\mathcal{A}(\Gamma_k)$ are labelled by $a \in \Gamma_k^*/\Gamma_k \cong \mathbb{Z}/2k\mathbb{Z}$, where Γ_k^* is the dual lattice of Γ_k

Another thing to notice about this simplified model is that the lattice chiral algebra depends only on k while the σ-model modulus in this case is R. Different factorisations of k into two coprime integers p and q will give different geometric realizations corresponding to a σ-model on a circle of radius square $R^2 = p/q$. We will encounter a similar feature when we study the case of $U_{m,K}$ algebra.

In this paper, we study the CFTs which arise from strings on complex multiplication tori and we propose their corresponding chiral algebra. We give a geometric interpretation for the $U_{m,K}$ algebra for any K and for $m = 2$ using the Gauss product [9]. We also show that the classification of the $U_{m,K}$ algebras in terms of rational points on a Grassmannian [10] is equivalent to tori of CM-type. We will make extensive use of the results in [6, 9, 10].

This paper is organized as follows. In Section 2, we introduce the $U_{m,K}$ matrix level Kac-Moody algebras and their modular invariant classification. In Section 3, we study complex multiplication tori and state the conditions for rationality drawing from the results in [8]. In Section 4, we formulate the problem of RCFTs based on complex multiplication tori in terms of rational points on Grassmannians and as such we relate it to the results in [10].

2. Matrix-level Kac-Moody algebras

In this section, we introduce the Kac-Moody algebras with matrix-valued level. We want to show that they constitute a consistent class of RCFTs by first describing their modular invariant partition functions. In Section 3 we relate them to σ-models on complex multiplication tori.

The abelian matrix level algebra $U_{m,K}$ with a matrix-values level K was studied in [10] where its modular invariant partition functions were classified. This algebra is a little different from the direct product form $U(1)_{k_1} \times \cdots \times U(1)_{k_m}$. That would correspond to a diagonal K matrix, where we consider the more general case, allowing K to be any integer-valued matrix.[1] For diagonal K, the geometric interpretation of $U_{m,K}$ is simply given by a torus $T^m = S_{k_1} \times \cdots \times S_{k_m}$, a product of circles with radii given in terms of the eigenvalues $k_1, \cdots, k_m$.

The algebra $U_{m,K}$ appears in the description of the fractional quantum Hall effect providing another motivation for its study. Recall the effective description of the fractional quantum Hall effect via a Chern-Simons theory with a gauge group $U(1)^m$ and a matrix valued level K [11, 12]. The Witten correspondence [13, 14] relates the RCFT based on a (non-twisted) affine Kac-Moody algebra (the Wess-Zumino-Witten model) and the Chern-Simons theory with corresponding gauge group G canonically quantized on a manifold $M = \Sigma \times \mathbb{R}$, where $\mathbb{R}$ is the time axis. For any compact gauge group G the Chern-Simons actions on a three-manifold are classified by an integer[2] $k \in \mathbb{Z}$. It was shown in [11] that Chern-Simons theories for the Abelian group $U(1)^m$ are classified by positive integer lattices with intersection forms K. Due to their potential application in the fractional quantum Hall effect, it is of interest to study the two-dimensional avatars of the $U(1)^m$ Chern-Simons theories based on matrix valued level K. This reduces to the study of two-dimensional RCFTs with the Kac-Moody algebra $U_{m,K}$.

Let Γ_K be a Euclidean, even, integral lattice of rank r with a positive definite integer-valued symmetric bilinear form

$$\Gamma_K = \mathbb{Z}\mathbf{e}_1 + \cdots + \mathbb{Z}\mathbf{e}_r, \quad \langle \mathbf{e}_i | \mathbf{e}_j \rangle = K_{ij} \in \mathbb{Z}. \tag{2.1}$$

The intersection matrix K_{ij} is a symmetric integral matrix with determinant $D < 0$. In the following we will refer to a lattice Γ_K and its intersection form K interchangeably with the hope that this will not cause any confusion.

The basis $\{\mathbf{e}_i\}$ of Γ_K is defined up to $GL(r,\mathbb{Z})$ transformations which preserve the determinant of K

$$\mathbf{e}_i \to A_i^j \mathbf{e}_j, \quad A \in GL(r,\mathbb{Z}), \quad \det A = \pm 1. \tag{2.2}$$

We will consider different matrix levels K_L for the holomorphic and K_R for the anti-holomorphic sectors which give rise to heterotic theories. The set of integrable representations of the affine algebra $U_{m,K}$ are labelled by $a \in P_+^{K_L} = \Gamma_{K_L}^*/\Gamma_{K_L}$ and $b \in P_+^{K_R} = \Gamma_{K_R}^*/\Gamma_{K_R}$ where $|K_L| = |K_R|$.

---

[1] We will also require that the diagonal elements of K to be even, $K_{ii} \in 2\mathbb{Z}$, since this is required for the invariance under the T transformation.

[2] If the three-manifold is a spin manifold then k can be half-integer [15].

We can associate with any integral lattice Γ_K of rank r a chiral vertex algebra $\mathcal{A}(\Gamma_K)$. There are r linearly independent $U(1)$ currents

$$J_i(z) = \sum_{n\in\mathbb{Z}} \frac{J_i^n}{z^{n+1}}. \tag{2.3}$$

They have the OPE

$$J_i(z)J_j(w) \sim \frac{K_{ij}}{(z-w)^2}. \tag{2.4}$$

In terms of the modes we have

$$\left[J_i^m, J_j^n\right] = mK_{ij}\delta_{m+n,0}. \tag{2.5}$$

These spin-one currents form a subalgebra $\mathcal{A}_r \subset \mathcal{A}(\Gamma)$ with rank r. The zero modes of the currents can be used to grade the Hilbert space into different superselection sectors with different charges $\mathbf{q} \in \Gamma_K$

$$\mathcal{H} = \oplus_{\mathbf{q}} \mathcal{H}_{\mathbf{q}}, \tag{2.6}$$

where

$$\mathcal{H}_{\mathbf{q}} = \left\{ |\mathbf{q}\rangle \ \mid \ J_i^0 |\mathbf{q}\rangle = q_i |\mathbf{q}\rangle \right\}. \tag{2.7}$$

The energy momentum tensor is given by the Sugawara construction

$$T(z) = \frac{1}{2} K^{ij} : J_i(z) J_j(z) :, \tag{2.8}$$

where $K^{ij} = K_{ij}^{-1}$ is the intersection form of the dual lattice.

From the OPE of $T(z)$ with itself we can read off the central charge

$$c = K^{ij} K_{ij} = r \tag{2.9}$$

The Virasoro generators are given by

$$L_n = \frac{1}{2} K^{ij} \sum_{m=-\infty}^{\infty} : J_i^{m+n} J_j^{-m} : \tag{2.10}$$

Now we specialize to rank two lattices Γ_K with basis $\mathbf{e}_1, \mathbf{e}_2$. Since the lattice is even, then its intersection form can be written as

$$K_{ij} = \langle \mathbf{e}_i | \mathbf{e}_j \rangle = \begin{pmatrix} 2a & b \\ b & 2c \end{pmatrix}. \tag{2.11}$$

We assume that $\gcd(a, b, c) = 1$ which corresponds to primitive quadratic forms.

The $GL(2, \mathbb{Z})$ transformation on the basis of Γ_K gives an equivalent lattice. The set of equivalence classes of primitive, even lattices is defined as

$$\mathcal{L}^p(D) := \left\{ \Gamma_K : D = -\det K \right\} / GL(2, \mathbb{Z}). \tag{2.12}$$

The RCFT data are given by $(\Gamma_{K_L}, \Gamma_{K_R}, \{\chi_a^{\Gamma_{K_L}}\}, \{\chi_a^{\Gamma_{K_R}}\})$ where $\chi_a^{\Gamma_{K_R}}$ are the characters which are proportional to the theta functions of the lattice

$$\theta_a^{\Gamma_{K_L}}(t) = \sum_{q\in\Gamma_{K_L}} q^{\frac{1}{2}(a+q)^2}. \tag{2.13}$$

We can compute the modular S matrix from transformation of the characters $(\chi_a^{\Gamma_{K_L}} = \theta_a^{\Gamma_{K_L}}/\eta^2)$ under $\tau \to -1/\tau$

$$\chi_a^{\Gamma_{K_L}}\left(\frac{-1}{\tau}\right) = \frac{\theta_a^{\Gamma_{K_L}}(-1/\tau)}{\eta(-1/\tau)^2}. \tag{2.14}$$

Using the transformation properties of the theta functions we find

$$S_{ab} = \frac{i}{\sqrt{D}} e^{-2i\pi(a\cdot b)}. \tag{2.15}$$

Now we can find the fusion rules using the Verlinde formula

$$\mathcal{N}_{ab}^{c} = \sum_d \frac{S_{ad}S_{bd}S_{cd}^{-1}}{S_{0d}} \tag{2.16}$$

The quantum dimension of a primary field labelled by a is

$$\frac{S_{a0}}{S_{00}} = 1, \tag{2.17}$$

i.e., all the primaries are simple currents. This simplifies the construction of all modular invariants of the algebra as was shown in [10].

The spectrum is encoded in the (genus-1) partition function

$$Z^{\Gamma_{K_L},\Gamma_{K_R}} = \sum_{a\in P_+^{K_L},\ b\in P_+^{K_R}} M_{a,b}\, \chi_a^{\Gamma_{K_L}} \overline{\chi_b^{\Gamma_{K_R}}}, \tag{2.18}$$

where the matrix $M_{a,b}$ is constrained to satisfy $M_{a,b} \in \mathbb{Z}_{\geq}$ and $M_{0,0} = 1$.

Modular invariance dictates that

$$SM = MS, \qquad TM = MT, \tag{2.19}$$

where the matrices S and T are unitary and symmetric and T is diagonal. They generate the modular group $SL(2,\mathbb{Z})$ and act on the characters as

$$\begin{aligned} S\cdot\chi_a &= \sum_{a,b\in P_+^K} S_{a,b}\chi_b, \\ T\cdot\chi_a &= \sum_{a,b\in P_+^K} S_{a,b}\chi_b. \end{aligned} \tag{2.20}$$

In [10], all modular invariants of the algebra $U_{m,K}$ were constructed in terms of even self-dual lattices Γ which contains Γ_{K_L} and Γ_{K_R}. In terms of the matrix K full modular invariance means $K_{ij} \in 2\mathbb{Z}$ which together with the fact that K is symmetric translates to an even integer lattice Γ_K.

The argument in [10] goes as follows: consider the heterotic partition function (2.18). Using $SM = MS$ and the unitarity of the S matrix we find

$$M_{a,b} = \sum S_{a,c} M_{c,d} S^*_{d,b} = \frac{1}{|K_L|} \sum \exp[2\pi i(b \cdot d - a \cdot c)] M_{c,d}. \tag{2.21}$$

Using the triangle inequality, the above equation becomes $|M_{a,b}| \leq |M_{0,0}| = 1$ with the equality iff

$$M_{c,d} \neq 0 \Longrightarrow a \cdot c = d \cdot b \pmod 1 \tag{2.22}$$

for all $c \in P_+^{K_L}, d \in P_+^{K_R}$. Define the set

$$\Omega = \bigcup_{a \in P_+^{K_L},\, b \in P_+^{K_R}} (a \oplus ib) + \left(\Gamma_{K_L} \oplus i\Gamma_{K_R}\right) \tag{2.23}$$

which is an even self dual lattice. The matrix $M_{a,b}$ in (2.18) satisfies

$$M_{a,b} = \begin{cases} 1 & \text{if } (a, ib) \in \Omega \\ 0 & \text{otherwise.} \end{cases} \tag{2.24}$$

Hence, the modular invariant partition functions of the $U_{m,K}$ current algebra are in one-to-one correspondence with the even self-dual lattices Γ which contain $(\Gamma_{K_L}; \Gamma_{K_R})$.

3. Strings on a complex multiplication torus

Given a complex number $\tau = \tau_1 + i\tau_2$ with $\tau_2 > 0$ we can construct an elliptic curve $E_\tau = \mathbb{C}/\Lambda$ where Λ is a lattice $\Lambda = (\mathbb{Z} + \tau\mathbb{Z})$ with τ the complex structure modulus of E_τ. The elliptic curve E_τ describes a torus which is given by identifying points in the complex plane related by the lattice translations generated by 1 and τ. The endomorphisms of E_τ are given by holomorphic maps $F : E_\tau \to E_\tau$. The only such holomorphic maps fixing 0 are induced by maps $G : \mathbb{C} \to \mathbb{C}$ of the form $G(z) = \lambda z$ where $\lambda \in \mathbb{C}$ is a constant such that $\lambda\Lambda \subseteq \Lambda$. Clearly any elliptic curve has trivial endomorphisms corresponding to multiplication by an integer $\lambda \in \mathbb{Z}$. We want to find the conditions on τ which gives non-trivial endomorphisms. The action of λ on the lattice Λ is represented on the generators as [7]

$$\begin{aligned} \lambda \cdot 1 &= m_1 \cdot 1 + n_1 \cdot \tau \\ \lambda \cdot \tau &= m_2 \cdot 1 + n_2 \cdot \tau, \end{aligned} \tag{3.1}$$

where $m_1, m_2, n_1, n_2 \in \mathbb{Z}$. By substituting the first equation in the second we learn that for an elliptic curve to have non-trivial endomorphisms, then τ must satisfy a quadratic equation with integer coefficients

$$a\tau^2 + b\tau + c = 0, \qquad \tau = \frac{-b + \sqrt{D}}{2a}, \quad D = b^2 - 4ac < 0. \tag{3.2}$$

Here we chose Im $\tau > 0$ since τ is the modular parameter of a torus.

The above equation means that τ takes values in the imaginary quadratic number field $\mathbb{Q}(D)$, the set of numbers of the form $\alpha + \beta\sqrt{D}$ with $\alpha, \beta \in \mathbb{Q}$.

Elliptic curves (or tori) for these special values of τ are said to have *complex multiplication* (or to be of CM-type). The σ-model on E_τ is specified by another parameter, the Kähler parameter ρ. The compactification of strings on E_τ is characterized by a momentum-winding Narain lattice, an even self-dual lattice $\Gamma(\tau, \rho)$ of rank 4, where the parameters τ and ρ live in the upper-half plane $\mathbb{H}^+$ subject to a group of discrete symmetries Ξ. The Narain moduli space of this compactification is [8]

$$\mathcal{M} = \frac{\mathbb{H}^+ \times \mathbb{H}^+}{\Xi}, \tag{3.3}$$

where Ξ is the group of discrete symmetries of E_τ

$$\Xi = PSL(2, \mathbb{Z})_\tau \times PSL(2, \mathbb{Z})_\rho \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2. \tag{3.4}$$

The first $\mathbb{Z}_2$ is a mirror symmetry which exchanges τ and ρ, i.e., $\mathbb{Z}_2 : (\tau, \rho) \mapsto (\rho, \tau)$. The second $\mathbb{Z}_2$ is a space-time parity transformation $\mathbb{Z}_2 : (\tau, \rho) \mapsto (-\bar{\tau}, -\bar{\rho})$, where the bar denotes complex conjugation. The last $\mathbb{Z}_2$ is a world-sheet orientation reversal $\mathbb{Z}_2 : (\tau, \rho) \mapsto (\tau, -\bar{\rho})$.

For generic values of τ and ρ, the conformal field theory is not rational and has an infinite number of primary fields. For special values of τ and ρ the infinite number of primary-fields representations reorganize into a finite set of representations of a bigger chiral algebra and the theory becomes rational. It was shown in [8] that CFTs based on E_τ are rational iff E_τ and its mirror are both of CM type, i.e., $\tau, \rho \in \mathbb{Q}(D)$ which implies that ρ, like τ, satisfies

$$a'\rho^2 + b'\rho + c' = 0, \qquad \rho = \frac{-b' + \sqrt{D'}}{2a'}, \quad D' = b'^2 - 4a'c' < 0. \tag{3.5}$$

Example of an RCFT which enjoys this property results when $\tau = \rho = e^{2\pi i/3}$. In this case the chiral vertex operator algebra is isomorphic to $SU(3)$ WZW model at level 1 [16]. Clearly, both τ and ρ satisfy

$$\tau^2 + \tau + 1 = 0. \tag{3.6}$$

In [8], the diagonal case was studied in detail where it was shown that the condition for a diagonal modular invariant is

$$\tau = fa\rho, \tag{3.7}$$

where $f \in \mathbb{Z}$ and $\tau, \rho \in \mathbb{Q}(D)$. The generalization to non-diagonal modular invariants was given in [9].

We can associate a quadratic form with the complex numbers τ and ρ. Write

$$Q(a,b,c) = \begin{pmatrix} 2a & b \\ b & 2c \end{pmatrix}, \qquad \tau_{Q(a,b,c)} = \frac{-b+\sqrt{D}}{2a} \tag{3.8}$$

where $D = b^2 - 4ac = -\det Q$ denotes the discriminant of the quadratic form Q. Q is called primitive if $\gcd(a,b,c) = 1$. The discriminant of the quadratic form Q is invariant under $SL(2,\mathbb{Z})$:

$$Q \longrightarrow S^t Q S, \quad S \in SL(2,\mathbb{Z}). \tag{3.9}$$

Now we can consider equivalence classes of quadratic forms under the action of $SL(2,\mathbb{Z})$, or more precisely $PSL(2,\mathbb{Z})$, since $S = \pm\mathbb{I}$ acts trivially. The set of equivalence classes is denoted by

$$Cl(D) = \left\{ Q(a,b,c) \,\middle|\, D = b^2 - 4ac < 0,\ a > 0 \right\} / \sim SL(2,\mathbb{Z}). \tag{3.10}$$

It is known that $Cl(D)$ is a finite set and we will denote the number of its elements by $h(D)$

$$Cl(D) = \left\{ \mathcal{C}_1, \dots, \mathcal{C}_{h(D)} \right\}. \tag{3.11}$$

Since $D < 0$, the complex number τ_Q lies in the upper-half plane $\mathbb{H}^+$. The $SL(2,\mathbb{Z})$ action on Q will induce a fractional linear transformation on τ_Q. The $PSL(2,\mathbb{Z})$ action on quadratic forms is compatible with the $PSL(2,\mathbb{Z})$ on $\mathbb{H}^+$. The $PSL(2,\mathbb{Z})$ orbits of $\tau_{Q(a,b,c)} \in \mathbb{H}^+$ depend on the class $\mathcal{C} = [Q(a,b,c)] \in Cl(D)$. Using the above mapping, then we can label the classes in $Cl(D)$ by points $[\tau_Q] \in \mathcal{F} = \mathbb{H}^+/PSL(2,\mathbb{Z})$. Since τ is the complex structure of a torus then fractional linear transformations on τ gives an equivalent torus. The classes $[\tau_Q]$ will give inequivalent tori in the Narain moduli space.

The same goes for ρ where we can also define equivalence classes of quadratic forms parametrized by points $[\rho_{Q'}] \in \mathcal{F} = \mathbb{H}^+/PSL(2,\mathbb{Z})$. The equivalence class of Narain lattices corresponding to an RCFT will be denoted by $\Gamma(\tau_{\mathcal{C}}, \rho_{\mathcal{C}'})$, where $\mathcal{C}$ and $\mathcal{C}'$ are the equivalence classes corresponding to τ and ρ.

Similarly, the group $GL(2,\mathbb{Z})$ acts on quadratic forms by the same formula as (3.9). We define the set of *improper* equivalence classes under $GL(2,\mathbb{Z})$ by

$$\tilde{Cl}(D) = \left\{ Q(a,b,c) \,\middle|\, D = b^2 - 4ac < 0,\ a > 0 \right\} / \sim GL(2,\mathbb{Z}). \tag{3.12}$$

We can associate with the lattice Γ_K a primitive quadratic form given by the intersection form K. Therefore, the equivalence classes of primitive lattices $[\Gamma_K]$ are in one-to-one correspondence with the equivalence classes of primitive, quadratic forms $[Q(a,b,c)]$. Hence, we can identify the set $\tilde{Cl}(D)$ with the set $\mathcal{L}^p(D)$ in (2.12).

Since $SL(2,\mathbb{Z}) \subset GL(2,\mathbb{Z})$ then $Cl(D) \subset \tilde{Cl}(D)$ and we have a surjection q

$$q : Cl(D) \to \tilde{Cl}(D), \quad \mathcal{C} \to \tilde{\mathcal{C}}, \tag{3.13}$$

where $q^{-1}(\tilde{\mathcal{C}})$ has either one or two classes.

We define the following projections of $\Gamma(\tau_{\mathcal{C}}, \rho_{\mathcal{C}'})$

$$\Pi_L := \Gamma(\tau_{\mathcal{C}}, \rho_{\mathcal{C}'}) \cap \mathbb{R}^{2,0}, \qquad \Pi_R := \Gamma(\tau_{\mathcal{C}}, \rho_{\mathcal{C}'}) \cap \mathbb{R}^{0,2}, \tag{3.14}$$

which correspond to the left and right momentum lattices, characterized by the vanishing of the left moving and right moving momenta, respectively.[3]

The modular invariant partition functions studied in [9] take the form:

$$\begin{aligned} Z^{\Pi_L,\Pi_R,\varphi}(q,\bar{q}) &= \frac{1}{|\eta(q)|^4} \sum_{a\in\Pi_L^*/\Pi_L} \theta_a^{\Pi_L}(q), \overline{\theta_{\varphi(a)}^{\Pi_R}(q)} \\ &= \sum_{a\in\Pi_L^*/\Pi_L} \chi_a^{\Pi_L}(q), \overline{\chi_{\varphi(a)}^{\Pi_R}(q)} \end{aligned} \tag{3.15}$$

where φ is a gluing map between the discriminant groups Π_L^*/Π_L and Π_R^*/Π_R. It satisfies $(\varphi(a), \varphi(b)) = (a, b)$, where $a \in \Pi_L^*/\Pi_L, b \in \Pi_R^*/\Pi_R$ and $(\cdot,\cdot)$ is the rational bilinear form on Π_L^*/Π_L which is induced from the bilinear form on Π_L. The theta function of the lattice Π_L is defined as

$$\theta_a^{\Pi_L}(q) = \sum_{v\in\Pi_L} q^{\frac{1}{2}(a+v)^2}. \tag{3.16}$$

The Gauss product

There is a binary operation which turns the set $Cl(D)$ into an abelian group: the Gauss product [9] takes two equivalence classes of quadratic forms and gives a third one with the same discriminant.

Let $\mathcal{C} = [Q_1(a_1,b_1,c_1)]$ and $\mathcal{C}' = [Q_2(a_2,b_2,c_2)]$ be two equivalence classes of quadratic forms with the same discriminant. We will restrict ourselves to primitive forms. We say that two quadratic forms $Q_1(a_1,b_1,c_1) \in \mathcal{C}$ and $Q_2(a_2,b_2,c_2) \in \mathcal{C}'$ are concordant if $a_1a_2 \neq 0$, $\gcd(a_1,a_2) = 1$ and $b_1 = b_2$. Then the Gauss product of $\mathcal{C}\star\mathcal{C}'$ is defined as

$$\big[Q_1\big(a_1,b,c_1\big)\big] \star \big[Q_2\big(a_2,b,c_2\big)\big] = \big[Q_3\big(a_3,b_3,c_3\big)\big], \tag{3.17}$$

where $a_3 = a_1a_2$, $b_3 = b$, and $c_3 = \frac{b^2-D}{4a_1a_2}$. It is important to mention that any pair of quadratic forms of the same discriminant can be $SL(2,\mathbb{Z})$ transformed to a concordant pair.

---

[3] More precisely it corresponds to an equivalence class of lattices.

The unit $\mathbf{1}_D$ of $Cl(D)$ with respect to the product $\star$ is represented by

$$\mathbf{1}_D = \begin{cases} [1, 0, -\frac{D}{4}]; & \text{if } D \equiv 0 \mod 4 \\ [1, 0, \frac{1-D}{4}]; & \text{if } D \equiv 1 \mod 4. \end{cases} \tag{3.18}$$

The quadratic form $Q_3(a_3, b_3, c_3)$ corresponds to a lattice with intersection form

$$Q_3\big(a_3, b_3, c_3\big) \equiv \begin{pmatrix} 2a_3 & b_3 \\ b_3 & 2c_3 \end{pmatrix}. \tag{3.19}$$

which gives an even integer lattice, an important fact which we will use when we construct the matrix levels K_L and K_R for the $U_{m,K}$ algebras.

The Gauss product is used to construct the intersection form[4] of the lattices Π_L and Π_R in terms of the equivalence classes $\mathcal{C}$ and $\mathcal{C}'$, corresponding to $\tau_{Q(a_1,b_1,c_1)}$ and $\rho_{Q(a_2,b_2,c_2)}$, respectively [9]:

$$\Pi_L = q(\mathcal{C} \star \mathcal{C}'^{-1}), \qquad \Pi_R = q(\mathcal{C} \star \mathcal{C}')(-1). \tag{3.20}$$

Here q is the natural map $Cl(D) \to \tilde{C}l(D)$ and $q(\mathcal{C} \star \mathcal{C}')(-1)$ means we multiply the quadratic form $q(\mathcal{C} \star \mathcal{C}')$ by -1.

As was shown in [9], to prove the above result one first constructs the $\mathbb{Z}$-basis for Π_R of $\Gamma(\tau_{\mathcal{C}}, \rho_{\mathcal{C}'})$ in terms of the equivalence classes of $\mathcal{C}$ and $\mathcal{C}'$ and then compares the resulting quadratic form of Π_R with $Q_3(a_3, b_3, c_3)$ and similarly for Π_L.

Now we have geometric data represented by the rational Narain lattice $\Gamma(\tau_{\mathcal{C}}, \rho_{\mathcal{C}'})$ which depends on the σ-model parameters τ and ρ (both τ and ρ are attached to an equivalence class of quadratic forms) and algebraic, RCFT data $(\Gamma_{K_L}, \Gamma_{K_R}, \{\chi_a\})$. The Gauss product can used to relate them in the following way. First we will look at K_L and K_R as quadratic forms and hence we can talk about their respective equivalence classes under the $SL(2, \mathbb{Z})$ action, as we did with Q. Now, a rational point in the Narain moduli space specified by the special values $\tau_{\mathcal{C}}$ and $\rho_{\mathcal{C}'}$ defines two equivalence classes of quadratic forms $\mathcal{C}$ and $\mathcal{C}'$. The mapping between the two sets of data is

$$K_L = q\big(\mathcal{C} \star C'^{-1}\big), \qquad K_R = q\big(\mathcal{C} \star \mathcal{C}'\big)(-1). \tag{3.21}$$

We propose using the algebra $U_{2,K}$ with matrix levels K_L and K_R as the chiral algebra for the RCFTs which arise from strings on CM-tori. To justify our claim, we notice that the characters of the RCFTs in [9] which are proportional to the theta functions (3.16) are the same as the characters (2.13) of the algebra $U_{2,K}$. Both sets of characters are based on lattices which are constructed from the same set of geometric data using the Gauss product. Also, the modular invariant partition functions (3.15) are a subset of the modular invariant partition functions (2.18) of the algebra $U_{2,K}$ for which

---

[4] This will define an equivalence class of lattices since the intersection form of a lattice define the basis up to $GL(2, \mathbb{Z})$ transformations.

$$M_{a,b} = \delta_{b,\phi(a)}. \tag{3.22}$$

The mapping (3.21) shows that the algebras $U_{m,K}$ can be given a geometric interpretation in terms of σ-models on complex multiplication tori.

We note that the above mapping is not one-to-one. We can't start from K_L and K_R and construct unique equivalence classes for τ and ρ. The matrices K_L and K_R are representatives of equivalence classes in the set $Cl(D)$ which have a finite number of elements. The algebraic description of the $U_{m,K}$ algebras depends only on K_L and K_R and as such is the same for all members of the set $Cl(D)$. On the other hand there is a geometric description for each member of the set $Cl(D)$. We conclude that the same $U_{m,K}$ algebra have many geometric avatars. This is similar to the case of a rational boson on a circle of raduis square $R^2 = p/q$ which is described a level $k = pq$ Kac-Moody $U(1)_k$ algebra. Starting from the algebra $U(1)_k$, there are many candidate rational boson theories, one for each factorization of k into two coprime integers $k = pq$.

4. Rational points on Grassmannians and CM tori

In this section we study the rationality conditions of a Narain lattice in more detail. We formulate the rationality in terms of rational points on a Grassmannian and we show that these points are equivalent to tori of complex multiplication type. Our argument will be based on the results in [6]. We will formulate some of these results in terms of rational points on a Grassmannian.

We consider a generic Narain lattice $\Gamma(\tau, \rho)$ of the σ-model on T^d/Λ with a B-field

$$\Gamma(\tau,\rho) = \Big\{ \big(P_L; P_R\big) := \frac{1}{\sqrt{2}}\big(\mu - \tilde{B}\lambda + \lambda; \mu - \tilde{B}\lambda - \lambda\big) \mid (\mu,\lambda) \in \Lambda^* \oplus \Lambda \Big\}, \tag{4.1}$$

where Λ^* and Λ are the momentum and winding lattices, respectively and $B = \Lambda^T \tilde{B} \Lambda$.

The holomorphic and anti-holomorphic vertex operators are characterized by $P_R = 0$ and $P_L = 0$ and they are parametrized by the values of their charges in Π_L and Π_R

$$\begin{aligned} \Pi_L &= \Big\{ \big(P_L; 0\big) := \frac{1}{\sqrt{2}}\big(\mu - \tilde{B}\lambda + \lambda; 0\big) | (\mu,\lambda) \in \Lambda^* \oplus \Lambda \Big\} \\ \Pi_R &= \Big\{ \big(0; P_R\big) := \frac{1}{\sqrt{2}}\big(0; \mu - \tilde{B}\lambda - \lambda\big) | (\mu,\lambda) \in \Lambda^* \oplus \Lambda \Big\}, \end{aligned} \tag{4.2}$$

We also define the following projections of the lattice $\Gamma(\tau, \rho)$

$$\begin{aligned} \widetilde{\Pi}_L &= \Big\{ \big(P_L; *\big) := \frac{1}{\sqrt{2}}\big(\mu - \tilde{B}\lambda + \lambda; *\big) | (\mu, \lambda) \in \Lambda^* \oplus \Lambda \Big\} \\ \widetilde{\Pi}_R &= \Big\{ \big(*; P_R\big) := \frac{1}{\sqrt{2}}\big(*; \mu - \tilde{B}\lambda - \lambda\big) | (\mu, \lambda) \in \Lambda^* \oplus \Lambda \Big\}, \end{aligned} \tag{4.3}$$

where the $*$ means we forget about the corresponding component of P.

Note that

$$\Pi_L \subseteq \widetilde{\Pi}_L, \qquad \Pi_R \subseteq \widetilde{\Pi}_R. \tag{4.4}$$

Since the lattice $\Gamma(\Lambda, B)$ is even, self-dual and integral then its straightforward to show that

$$\Pi_L^* \cong \widetilde{\Pi}_L, \qquad \Pi_R^* \cong \widetilde{\Pi}_R, \tag{4.5}$$

i.e., $\widetilde{\Pi}_L$ is the dual of Π_L and the same for $\widetilde{\Pi}_R$.

Rationality can be expressed in terms of the rank of Π_L and Π_R . The Narain lattice $\Gamma(\tau, \rho)$ is rational if and only if [6]

$$\text{rank}\big(\Pi_L\big) = \text{rank}\big(\Pi_R\big) = d. \tag{4.6}$$

To see how the above conditions lead to rationality we recall the operator product expansion of the vertex operators take the form

$$V[P](z, w) \times V[P'](\bar{z}, \bar{w}) \sim c_{PP'}(z - w)^{P_L P_L'}(\bar{z} - \bar{w})^{P_R P_R'} V[P + P']. \tag{4.7}$$

On the circle $(z - w)(\bar{z} - w) = 1$, the above OPE takes the form

$$V[P](z, w) \times V[P'](\bar{z}, \bar{w}) \sim c_{PP'}(z - w)^{P_L P_L' - P_R P_R'} V[P + P'] \tag{4.8}$$

which defines the natural metric on the lattice of charges $\Gamma(\tau, \rho)$ to be $P \circ P' = P_L P_L' - P_R P_R$.

RCFTs are characterized by the appearance of extra holomorphic vertex operators which extend the chiral algebra. For a generic Narain lattice $\Gamma(\tau, \rho)$, the only holomorphic vertex operator is the one corresponding to the vacuum $P_L = P_R = 0$. Since any field is mutually local[5] to the vacuum, then the set of allowed representations $\mathcal{V} = \{V[P], P \in \Gamma(\tau, \rho)\}$ is infinite. However extra holomorphic vertex operators $\mathcal{W}[P]$ appear for $p \in \Pi_L$. The requirement of locality with respect to $\mathcal{W}[P]$ restricts the set of allowed irreducible representation of the above OPE to be $a \in \Pi_L^*/\Pi_L$, where Π_L^* is the lattice dual to Π_L so it contains charges which have integer $\circ$ product with Π_L. The condition for rationality translates to the requirement that Π_L be a finite index sublattice of Π_L^* so that the set Π_L^*/Π_L has a finite cadinality. This happens when Π_L have a finite rank which is the condition in (4.6).

---

[5] Two fields are mutually local if their OPE doesn't contain branch cuts. In other words the $\circ$ product of their charges is integer.

The Narain moduli space of conformal field theories is isomorphic to the moduli space of even, self-dual lattices with signature (d, d)

$$\mathcal{M}_d = O(\Gamma^{d,d})\backslash O(d,d)/(O(d) \times O(d)), \tag{4.9}$$

where $\Gamma^{d,d}$ denotes the standard even self-dual lattice of signature (d, d) and $O(\Gamma^{d,d})$ its automorphism group. In the special case of $d = 2$ this gives the moduli space in (3.3). The above moduli space is also the Grassmannian of space-like d-planes in $\mathbb{R}^{d,d}$. The modular invariant partition functions in [10] are classified using rational points on the Grassmannian (4.9).

Note that

$$P_L \in W = \Pi_L \otimes \mathbb{R}, \qquad P_R \in W^{\perp} = \Pi_R \otimes \mathbb{R}, \tag{4.10}$$

where W is a space-like d-plane which correspond to a point on the Grassmannian (4.9) and $W^{\perp}$ is its orthogonal complement in $\mathbb{R}^{d,d}$.

The sets Π_L and Π_R in general are not lattices, since the set of vectors which span Π_L and Π_R will not remain linearly independent over $\mathbb{Z}$ when restricted to W and $W^{\perp}$. However, if W is a *rational* point on the Grassmannian (4.9) then Π_L and Π_R become lattices of rank d. A rational point on the Grassmannian (4.9) is a subspace W with basis $\{f_m\}$ which can be written over $\mathbb{Q}$ in terms of the preferred orthonormal basis e_i of $\mathbb{R}^{d,d}$

$$f_m = Q_{mi} e_i, \quad Q_{mi} \in \mathbb{Q}. \tag{4.11}$$

The above equation implies that the basis of W are rational vectors

$$\langle f_m | f_n \rangle \in \mathbb{Q}. \tag{4.12}$$

Since any group generated by rational vectors is a lattice, i.e., it can be generated by linearly independent vectors over $\mathbb{Z}$. Then Π_L which is the $\mathbb{Z}$-span of f_m is a lattice of rank $d = \dim(W)$ and the same for Π_R. It was shown in [6] that for $d = 2$

$$\mathrm{rank}\big(\Pi_L\big) = \mathrm{rank}\big(\Pi_R\big) = 2 \longleftrightarrow \tau, \rho \in \mathbb{Q}(D). \tag{4.13}$$

which in our case implies that rational points on the Grassmannian (4.9) correspond to complex multiplication tori. This is another way to see that RCFTs constitute a dense subset of the set of CFTs, since the set of rational points on a Grassmannian is dense which is consistent with the findings in [8] that the values $\tau, \rho \in \mathbb{Q}(D)$ corresponding to RCFTs are dense in $\mathbb{H}^+ \times \mathbb{H}^+$.

5. Conclusion

We studied the RCFTs which arise from strings on complex multiplication tori corresponding to specific values of the moduli $\tau, \rho \in \mathbb{Q}(D)$ inside the Narain moduli space. We proposed using the matrix level algebra $U_{2,K}$ as the chiral algebra for these RCFTs. The Gauss product was used to construct the levels K_L and K_R out of the geometric data represented by τ and ρ. This also gives a geometric interpretation to the algebra $U_{2,K}$. By formulating the problem is terms of rational points on a Grassmannian we showed that the set of RCFTs is a dense subset in the Narain moduli space. This agrees with the observation in [8] that the values of $\tau, \rho \in \mathbb{Q}(D)$ which produce RCFTs are dense in the Narain moduli space.

The Gukov-Vafa criterion characterizes the rationality of the CFT in terms of the geometry of the target space. In the Gannon classification, rationality is derived from the algebraic properties of the chiral algebra without reference to any target space interpretation. We showed that the geometric and the algebraic characterization of rationality are related and that the density property of RCFTs can be seen in both pictures.

It would be interesting to generalize this to d-dimensional tori and also to general $U_{d,K}$ algebras. Another generalization would be the study of WZW models based on matrix level affine Kac-Moody algebras for non-Abelian groups. This would also be motivated by their relevance to a similar problem for Chern-Simon theories.

References

[1] P. Di Francesco, P. Mathieu, and D. Sénéchal, *Conformal field theory*, .

[2] P. H. Ginsparg, *Applied Conformal Field Theory*, `hep-th/9108028`.

[3] J. L. Cardy, *Operator content of two-dimensional conformally invariant theories*, *Nucl.Phys.* **B270** (1986) 186–204.

[4] D. Gepner, *Space-time supersymmetry in compactified string theory and superconformal models*, *Nucl.Phys.* **B296** (1988) 757.

[5] J. A. Harvey, G. W. Moore, and C. Vafa, *Quasicrystalline compactification*, *Nucl.Phys.* **B304** (1988) 269–290.

[6] K. Wendland, *Moduli spaces of unitary conformal field theories*, . PhD Thesis.

[7] G. W. Moore, *Arithmetic and attractors*, `hep-th/9807087`.

[8] S. Gukov and C. Vafa, *Rational conformal field theories and complex multiplication*, *Commun.Math.Phys.* **246** (2004) 181–210, [`hep-th/0203213`].

[9] S. Hosono, B. H. Lian, K. Oguiso, and S.-T. Yau, *Classification of $c = 2$ rational conformal field theories via the Gauss product*, *Commun.Math.Phys.* **241** (2003) 245–286, [`hep-th/0211230`].

[10] T. Gannon, *U(1)-m modular invariants, N=2 minimal models, and the quantum Hall effect*, *Nucl.Phys.* **B491** (1997) 659–688, [`hep-th/9608063`].

[11] D. Belov and G. W. Moore, *Classification of Abelian spin Chern-Simons theories*, `hep-th/0505235`.

[12] A. Cappelli and G. R. Zemba, *Modular invariant partition functions in the quantum Hall effect*, *Nucl.Phys.* **B490** (1997) 595–632, [`hep-th/9605127`].

[13] E. Witten, *Quantum field theory and the Jones polynomial*, *Commun.Math.Phys.* **121** (1989) 351.

[14] G. W. Moore and N. Seiberg, *Taming the conformal zoo*, *Phys.Lett.* **B220** (1989) 422.

[15] R. Dijkgraaf and E. Witten, *Topological gauge theories and group cohomology*, *Commun.Math.Phys.* **129** (1990) 393.

[16] C. Vafa, *Quantum symmetries of string vacua*, *Mod.Phys.Lett.* **A4** (1989) 1615.